

& in D. C. Prodr. xii (1848) 177. *L. macrophyllus*, Benth. ll. cc. (1833, 1848). *L. virginicus*, var. *macrophyllus*, Gray, Proc. Am. Acad. viii. (1870) 285 & Syn. Fl. ii. pt. 1, 353. *L. communis*, Bicknell in Britton, Man. (1901) 803; Small, Fl. 1048. *L. membranacea*, Bicknell, l. c. (1901) 804.

GRAY HERBARIUM.

ADDENDA TO THE FLORA OF VERMONT.

W. W. EGGLESTON (revised by E. BRAINERD).

SINCE the publication of the Vermont Flora¹ in 1900 there has been a marked increase of interest in botanical field-work in various localities of the State, resulting in the discovery of many additional species and varieties. At the same time several of the more difficult genera of the Northeastern United States have been diligently worked over by specialists, and many old species have been divided into two, or several, or many species. Though in some cases the status of the proposed species is as yet somewhat uncertain, they have nevertheless been admitted with considerable freedom into this provisional list as plants that challenge for the time being the observation and criticism of the field botanist.

It is expected that within two years a new edition of the Vermont Flora will be published, and any further additions or corrections will be most gratefully appreciated; they may be communicated to the writer, or to the President of the Vermont Botanical Club, Ezra Brainerd of Middlebury, or to its Secretary, L. R. Jones of Burlington.

INDIGENOUS OR WELL-ESTABLISHED PLANTS.²

Aspidium acrostichoides, Swartz, var. *incisum*, Gray. Occasional. A form of this with crisp edges also occurs: Greensboro, *Miss H. M. Hodge*; Burlington, *Mrs. F. A. Richardson*.

¹ Brainerd, Jones & Eggleson, Flora of Vermont (Contributions to the Botany of Vermont, viii). Burlington, 15 December, 1900.

² As in the Flora of Vermont, the names of plants indigenous in North America are printed in full face type and those of species introduced from the Old World in small capitals.

- A. simulatum**, Davenport. Brattleboro, *Mrs. F. B. Horton*.
- A. spinulosum** × **marginale**. (*Dryopteris Pittsfordensis*, Slosson, *RHODORA*, vi. 75.) Pittsford, *Miss Margaret Slosson*.
- Asplenium ebeneum**, Ait. var. **Hortonae**, Davenport, *RHODORA*, iii. 2, plate 22. Brattleboro, *Mrs. F. B. Horton*; Pittsford, *Miss K. A. French*.
- Osmunda Claytoniana**, L. var. **dubia**, A. J. Grout. An abnormal form, with pinnules separated by a distance of one-half to two-thirds their breadth, the upper ones greatly elongated and pinnately cleft. Newfane, *Grout*. (See Papers of Linn. Fern Chapter, August, 1898.)
- Pteris aquilina**, L. var. **pseudocaudata**, Clute. (*P. latiuscula*, Desv.) Manchester, *F. A. Ross*,—the only New England station.
- Equisetum pratense**, Ehrh. Newark, *Kennedy*.
- Lycopodium clavatum**, L. var. **monostachyon**, Greville & Hooker. (See *RHODORA*, iii. 237, and iv. 57, 100.) Westmore, *Kennedy*; Windham, Westminster, Putney, *W. H. Blanchard*; Ira, *D. S. Carpenter*.
- L. complanatum**, L. var. **Wibbei**, Harberer, *RHODORA*, vi. 102. Westmore, *Kennedy*.
- Isoetes macrospora**, Dur. Willoughby Lake, *Kennedy*.
- I. Tuckermani**, A. Br. var. **borealis**, A. A. Eaton. Grout Pond, Stratton, *Eggleson and Grout*.
- Calamagrostis hyperborea**, Lange, var. **elongata**, Kearney. Willoughby Mt., *E. & C. E. Faxon*, 1887; specimen in Gray Herb.
- Lolium temulentum**, L. Burlington, *Jones*.
- PANICUM MILIACEUM**, L. Lumber yard, Burlington, *Brainerd*.
- P. minus**, Nash. Head of Lake Memphramagog, *E. Tuckerman*, 1859. (See *RHODORA*, iii. 105.)
- P. sphaerocarpon**, Ell. Railroad, Putney, *W. H. Blanchard*.
- Carex aenea**, Fernald, Proc. Am. Acad. xxxvii. 480 (1902). Previously confused with *C. foenea*, Willd. Cliffs and sand plains in Rutland and Addison Counties.
- C. aestivalis**, M. A. Curtis. Pownal, *J. R. Churchill*.
- C. canescens**, L. var. **disjuncta**, Fernald, l. c. 488. The common form; typical *C. canescens* has been recognized in Vermont only from Ripton.
- C. canescens**, var. **subloliacea**, Laestad. Cold bogs, Mt. Mansfield, *Brainerd*; Mud Pond, Wallingford, *Brainerd and Eggleson*.
- C. Crawfordii**, Fernald, l. c. 469. (*C. scoparia*, var. *minor*, Boott, in part.) Cold sterile soil; frequent.
- C. Crawfordii**, var. **vigens**, Fernald. Middlebury, *Brainerd*; Rutland, *Eggleson*; St. Johnsbury, *Hazen*.
- C. hystricina**, Muhl. var. **Dudleyi**, Bailey. Manchester, *Miss M. A. Day*.
- C. intumescens**, Rudge, var. **Fernaldii**, Bailey. St. Johnsbury, *Hazen*; Middlebury, *Brainerd*.
- C. mirabilis**, Dewey, var. **perlonga**, Fernald, l. c. 473. Little Notch, Bristol, *Brainerd*.
- C. mirabilis**, var. **tincta**, Fernald, l. c. 473. St. Johnsbury, *Hazen*.
- C. Pennsylvanica**, Lam. var. **lucorum**, Fernald, l. c. 505. Middlebury and Burlington, *Brainerd*; Pownal, *J. R. Churchill*; Fair Haven, *Eggleson*.

- C. scoparia*, Schkuhr, var. *moniliformis*, Tuckerm. Burlington, *Hazen*; Knight's Island, *Brainerd*.
- C. scoparia*, var. *condensa*, Fernald, l. c. 486. West Rutland, *Eggleson*; Willoughby Lake, *E. F. Williams*.
- C. setacea*, Dewey. Sudbury, *Brainerd*.
- C. straminea*, Willd. var. *echinodes*, Fernald, l. c. 474. Dame Island, North Hero, *Brainerd*.
- C. umbellata*, Schkuhr, var. *brevirostris*, Boott. (See Proc. Am. Acad. xxxvii. 507.) Snake Mt., *Brainerd*.
- C. varia*, Muhl. Pownal, *Eggleson*; Mallett's Head, Colchester, *Brainerd*; Burlington, *Pringle*.
- C. vesicaria*, var. *distenta*, Fries. (See RHODORA, iii. 54.) Knight's Island, *Brainerd*.
- C. vesicaria*, var. *jejuna*, Fernald, RHODORA, iii. 53. Common.
- C. gracillima* × *castanea*, (?). St. Johnsbury, *Hazen*.
- [*C. scabrata* × *crinita*, Fernald. By an error this hybrid was credited to Vermont in Mr. Fernald's check-list of New England Carices (RHODORA, iv. 222.) The plant was from Mt. Clinton, N. H.]
- Scirpus heterochaetus*, Agnes Chase, RHODORA, vi. 70, plate 53, d. Lake Champlain, Milton, *A. J. Grout*, 1893; the only station known, as yet, in New England for this new species of bulrush.
- S. lineatus*, Michx. Roadside in moist clay soil, Middlebury, *Brainerd*; Bristol, *A. C. Dike*.
- Arisaema stewardsonii*, Britton, Manual, p. 1045. Distinguished by its green, fluted spathe. Hartland, Pownal, Ferrisburg, *Eggleson*.
- Juncus alpinus*, Vill. Willoughby Mt. and in bog near hotel, *Eggleson*.
- J. articulatus*, L. var. *obtusatus*, Engelm. (See RHODORA, vi. 35.) Manchester, *Miss M. A. Day*.
- J. secundus*, Beauv. Twin Mts., West Rutland, *Eggleson*.
- J. tenuis*, Willd. var. *Williamsii*, Fernald, RHODORA, iii. 60, plate 23. "Sandfield," Rutland, *Eggleson*; Willoughby Lake, *Kennedy*.
- ALLIUM SCHOENOPRASUM*, L. Willoughby Lake, *Kennedy*. The plants reported in the Vermont Flora are var. *Sibiricum*, Hartm.
- Convallaria majalis*, L. Thickly covering an area of about six square rods in a mucky bog, remote from dwellings, Waltham, *Miss Ruth B. Fisher*.
- Habenaria psychodes* × *lacera*, Andrews, RHODORA, iii. 245 (cf. v. 263). Pownal, *Andrews and White*.
- Listera auriculata*, Wiegand. Lyndon, *J. W. Congdon*.
- Pogonia affinis*, Austin. One plant, Burlington, *Mrs. Henry Holt*. (See RHODORA, iv. 216.)
- Populus balsamifera*, L. var. *candidans*, Gray. Persists about old homesteads.
- SALIX ALBA*, L. var. *COERULEA*, Koch. "Planted and spontaneous, Barre," *Miss Eunice D. Smith*.
- HUMULUS JAPONICA*, Sieb. & Zucc. Waste places, Brattleboro, *Miss Anne S. Angell*; Burlington, *Jones*; Bellows Falls, *W. H. Blanchard*.

Polygonella articulata, Meisn. "Roadside, Vernon road, Brattleboro, Aug. 17, 1844," *T. W. Higginson*. Specimen, now in Gray Herb., was found pressed beside the above note in Col. Higginson's copy of Bigelow's *Florula Bostoniensis*. Colchester Point, *Pringle*.

POLYGONUM AVICULARE, L. var. **VEGETUM**, Ledeb. (See *RHODORA*, iv. 70.) Frequent in rich shaded soil. This was by error listed in the Vermont Flora as *P. ramosissimum*, Michx., which name should therefore be omitted.

Rumex salicifolius, Weinm. Fort Ethan Allen, *L. B. Sprague*.

CHENOPodium ALBUM, L. var. **VIRIDE**, Moq. Common.

LYCHNIS FLOS-CUCULI, L. Weed, roadsides and meadows, Craftsbury, *Jones*.

Silene stellata, Ait. Mr. Fernald kindly reports that this species is represented in the Michaux Herbarium at the Jardin des Plantes in Paris by a characteristic specimen labelled by Michaux, "Vermont et Nouvelle Angleterre."

Actaea neglecta, Gilman. (*A. eburnea*, Rydb.) Willoughby Mt., *H. H. Rusby*. This plant has been passing as white-fruited *A. spicata*, var. *rubra*, Ait. It is perhaps a hybrid of this with *A. alba*, Bigel.

Ranunculus Alleghaniensis, Britton. Smuggler's Notch, *E. F. Williams*.

R. sceleratus, L. Colchester, *Mrs. Flynn*.

Arabis brachycarpa, Britton. (See *RHODORA*, v. 225.) Sandy and rocky shores of Lake Champlain and vicinity. This and the following species replace *A. confinis*, Watson.

A. Drummondi, Gray. Rocky hills, western Vermont.

Sisymbrium canescens, Nutt. (*Sophia brachycarpa*, Rydb.) Hogback Island, Colchester, *Mrs. Flynn*.

Crataegus anomala, Sarg. *RHODORA*, iii. 74. Bridport, Clarendon. This and the following proposed species of this genus have been collected principally by Brainerd and Eggleston in western Vermont, and by W. H. Blanchard in Windham Co.

C. ascendens, Sarg. *RHODORA*, v. 141. Middlebury, Bristol, New Haven; West Rutland; Westminster, Putney.

C. asperifolia, Sarg. *RHODORA*, iii. 31. New Haven, Middlebury; Bellows Falls.

C. Brainerdi, Sarg. *RHODORA*, iii. 27. Middlebury, New Haven; West Rutland; Bellows Falls, Westminster; St. Johnsbury, *Miss I. M. Paddock*.

C. coccinea, L. Marsh Hill, Ferrisburg; Windham.

C. Champlainensis, Sarg. *RHODORA*, iii. 20. Frequent in the Champlain valley.

C. contigua, Sarg. *RHODORA*, v. 115. Stockbridge, *Eggleston*.

C. delucida, Sarg. *RHODORA*, v. 139. Widely distributed throughout the State.

C. demissa, Sarg. *RHODORA*, v. 139. Charlotte, *Horsford*.

C. dilatata, Sarg. *Bot. Gaz.* xxxi. 9. Frequent in the Champlain valley.

C. dissona, Sarg. *RHODORA*, v. 60. Near Bellows Falls.

C. Egglestoni, Sarg. *RHODORA*, iii. 30. One of the most common species, especially of higher altitudes up to 2300 ft.

- C. exclusa*, Sarg. *RHODORA*, v. 108. Bald Mt. Shrewsbury.
- C. flabellata*, Spach. (See *RHODORA*, iii. 75.) Bellows Falls, Westminster, Putney.
- C. genialis*, Sarg. *RHODORA*, v. 148. Common.
- C. glaucophylla*, Sarg. *RHODORA*, v. 140. Bennington, Arlington; Bellows Falls, Westminster, Putney.
- C. Gravesii*, Sarg. *RHODORA*, v. 159. Vergennes, New Haven.
- C. Holmesiana*, Ashe. (See *Bot. Gaz.* xxxi. 10.) Common.
- C. intricata*, Lange. Twin Mts., West Rutland; Arlington.
- C. Jesupi*, Sarg. *RHODORA*, v. 61. Twin Mts., West Rutland.
- C. lobulata*, Sarg. *RHODORA*, iii. 22. Common in Champlain valley.
- C. matura*, Sarg. *RHODORA*, iii. 24 & v. 144. Common.
- C. media*, Sarg. *RHODORA*, v. 150. Summit of Willoughby Mt., *E. Faxon*.
- C. membranacea*, Sarg. *RHODORA*, v. 186. Middlebury.
- C. modesta*, Sarg. *RHODORA*, iii. 28. Twin Mts., West Rutland; Arlington.
- C. pentandra*, Sarg. *RHODORA*, iii. 25. Common.
- C. polita*, Sarg. *RHODORA*, v. 111. Bennington, Pownal; Cornwall; Bellows Falls, Westminster.
- C. praecoqua*, Sarg. *RHODORA*, v. 167. Bellows Falls, Westminster, Putney; Lemington.
- C. Pringlei*, Sarg. *RHODORA*, iii. 21. Common.
- C. punctata*, Jacq. var. *aurea*, Ait. Common.
- C. scabrida*, Sarg. *RHODORA*, iii. 29. Common.
- C. submollis*, Sarg. *Bot. Gaz.* xxxi. 7. Weybridge Monument.
- C. succulenta*, Link. Middlebury, New Haven.
- Potentilla Canadensis*, L. West Rutland, *Egglesston*.
- Pyrus arbutifolia*, L. f. Westminster, Bellows Falls, *W. H. Blanchard*.
- P. communis*, L. Frequently spontaneous.
- Rosa Gallica*, L. Established in many places, *W. H. Blanchard*.
- Rubus nigricans*, Rydb. Common in the mountain towns.
- R. occidentalis*, L. A form with amber-colored fruit. Burlington, *L. P. Sprague*; Weathersfield, *W. H. Blanchard*.
- Desmodium rotundifolium*, DC. Dummerston, *L. Wild*; Bellows Falls, *W. H. Blanchard*.
- Lathyrus tuberosus*, L. Vergennes, *Miss Ruth B. Fisher*. (See *RHODORA*, iii. 230.)
- Lespedeza Nuttallii*, Darl. Vernon, *W. H. Blanchard*.
- L. Stuvei*, Nutt. Vernon, *W. H. Blanchard*.
- Linum medium*, Greene. Arlington, *W. H. Blanchard*.
- L. sulcatum*, Riddell. Pownal, *W. H. Blanchard*.
- Ampelopsis quinquefolia*, Michx. var. *muronum*, Rehder. Red Rocks, Burlington, *Rehder and Jones*.
- Lechea minor*, L. Vernon, *W. H. Blanchard*.
- Viola cucullata*, Ait. Springs, cold brooks and boggy meadows; common. (For an account of this and the following species of *Viola* see *RHODORA*, vi. 8-17.)
- V. latiuscula*, Greene, Pitt. v. 93. Dry sterile soil; common.

- V. papilionacea**, Pursh. Meadows and orchards; occasional.
- V. septentrionalis**, Greene. Pitt. iii. 334. Groves, hill pastures, especially under young conifers; frequent.
- V. sororia**, Willd. Rich open woods, moist meadows and pastures; common.
- V. venustula**, Greene, Pitt. iii. 335. Moist woods, and boggy meadows; frequent. (Hardly distinct from *V. affinis*, LeConte.)
- Gaylussacia resinosa**, Torr. & Gray, var. *glaucocarpa*, Robinson, RHODORA, ii. 83. Westminster, Pownal, *W. H. Blanchard*.
- Pyrola asarifolia**, Michx. Norwich, *Eggleson*; Westmore, *Kennedy*.
- Vaccinium Oxycoccus**, L. var. *intermedium*, Gray. (See RHODORA, iv. 237, plate 40.) Colchester, *Oakes*; Willoughby Lake, *Kennedy*.
- ANAGALLIS ARVENSIS**, L. Lawns, Bradford, *Miss A. E. Bacon*.
- A. ARVENSIS**, L. var. *CAERULEA*, Ledeb. Bradford, *Miss Bacon*. (See RHODORA, iv. 185.)
- Apocynum medium**, Greene. Willoughby Lake, *Kennedy*.
- Phlox subulata**, L. Old cemeteries; St. Johnsbury, *Miss I. M. Paddock*; Putney, Stratton, *W. H. Blanchard*.
- Myosotis arvensis**, Lam. Sharon, *Dr. Baxter*; Townshend, *W. H. Blanchard*.
- SYMPHYTUM ASPERRIMUM**, Sims. Dummerston, Vernon, *W. H. Blanchard*; Stockbridge, Rutland, *Eggleson*.
- Verbena angustifolia**, Michx. Arlington, *W. H. Blanchard*.
- V. hastata** × *urticifolia*. Pownal, *W. H. Blanchard*.
- CALAMINTHA ACINOS**, L. Townshend, *W. H. Blanchard*.
- Hedoma hispida**, Pursh. Burlington, *F. H. Hollister*.
- LAMIUM ALBUM**. Burlington, *Mrs. Flynn*.
- Lycopus uniflorus**, Michx. (*L. communis*, Bicknell & *L. membranacea*, Bicknell, Britton Manual p. 803.) Wet meadows and thickets; common. (See RHODORA, vi. 134.)
- Solanum rostratum**, Dunal. Barre, *Miss Eunice D. Smith*.
- S. Carolinense**, L. Woodstock, *L. Wild*; Westminster, *W. H. Blanchard*.
- VERONICA CHAMAEDRYS**, L. Glebe Mt., Windham, *W. H. Blanchard*.
- V. serpyllifolia**, L. var. *borealis*, Laest. (See RHODORA iv. 193.) Peacham, *F. Blanchard*; Willoughby Lake, *Kennedy*; St. Johnsbury, *Miss I. M. Paddock*.
- Utricularia clandestina**, Nutt. Willoughby Lake, *Kennedy*.
- Galium boreale**, L. Burlington, *Mrs. Flynn*; Manchester, *W. H. Blanchard*.
- G. SYLVATICUM**, L. Cultivated, but also well established on a shady ledge, Orwell, *Brainerd*; Peacham, *F. Blanchard*.
- Campanula uliginosa**, Rydb. Britton Manual 885. Separated from *C. aparinoides*; apparently more common.
- Lobelia cardinalis**, L. A form with white flowers. St. Albans, *M. D. Merrick*.
- ANTHEMIS ARVENSIS**, L. Westminster, Townshend, *W. H. Blanchard*.
- Aster Lindleyanus**, Torr. & Gray. var. *comatus*, Fernald, n. var. "Stems and under surfaces of the midribs copiously pubescent with soft white hairs.—NEW BRUNSWICK, dry woods near Four Falls, Grand Falls Par-

ish, August 14, 1901, *M. L. Fernald*; MAINE, river thicket, Fort Fairfield, September 19, 1900 — type — and sandy river terrace, Fort Fairfield, September 27, 1901, *M. L. Fernald*; VERMONT, Ripton, August 26, 1903, and Bellows Falls, August 24, 1903, *W. H. Blanchard*: SASKATCHEWAN, without locality, 1858, *Bourgeau*. The Vermont plants are less extreme than those from New Brunswick, Maine and Saskatchewan, but are evidently close to them."

A. *longifolius*, Lam. Willoughby Lake, *Kennedy*; Pownal, *Eggleson*; Connecticut valley, *W. H. Blanchard*.

A. *longifolius*, var. *villicaulis*, Gray. Connecticut valley, *W. H. Blanchard*.

A. *paniculatus*, Lam. var. *acutidens*, Burgess. Westminster, *W. H. Blanchard*.

A. *paniculatus*, var. *cinerascens*, Fernald, *RHODORA*, i. 188. Westminster, *W. H. Blanchard*.

A. *paniculatus*, var. *lanatus*, Fernald, *RHODORA*, vi. 131. Willoughby Lake, *Kennedy*.

A. *paniculatus*, var. *simplex*, Burgess. Westminster, *W. H. Blanchard*.

A. *Schreberi*, Nees. Common in southern Vermont, *W. H. Blanchard*.

A. *tardiflorus*, L. var. *vestitus*, Fernald, *RHODORA*, i. 188. Westminster, *W. H. Blanchard*.

Bidens vulgata, Greene. Previously confused with *B. frondosa*. Common.

Eupatorium perfoliatum, L. forma *purpureum*, Britton. Putney, *W. H. Blanchard*.

Hieracium Marianum, Willd. Bellows Falls, *W. H. Blanchard*.

H. PILOSELLA, L. St. Johnsbury, *Miss I. M. Paddock*.

Senecio Balsamitae, Muhl. var. *praelongus*, Greenman, *RHODORA*, iii. 6. Manchester, *Miss Day*.

Silphium perfoliatum, L. Burlington, *Jones*; Westminster, *W. H. Blanchard*.

Solidago juncea, Ait. var. *scabrella*, Gray. Manchester, *W. H. Blanchard*.

S. odora, Ait. Vernon, *W. H. Blanchard*. Also at Walpole, N. H.

TARAXACUM OFFICINALE, Weber, var. *PALUSTRE*, Blytt. (See *RHODORA*, iv. 155.) Rutland, *Eggleson*.

Xanthium Pennsylvanicum Wallr. (*X. commune*, Britton.) Bellows Falls, *Brainerd*; Burlington, *Jones*.

X. speciosum, Kearney. Burlington, *Jones*.

CASUAL PLANTS.

The following is a list of plants that have recently appeared spontaneously in cultivated fields, along railroads or near buildings. They are either garden escapes or waifs coming in mostly with foreign seed and should not, as yet, be regarded as having a proper place in our Flora.

Juncus Gerardi, Loisel. On the margin of a marshy meadow very near the railway station at New Haven Junction; perhaps from saltmarsh hay often used in packing. *Brainerd*.

ORNITHOGALUM UMBELLATUM, L. Brattleboro, *W. H. Blanchard*.

IRIS PSEUDACORUS, L. Burlington, *F. A. Ross*.

- RUMEX MONTANUM, Desf. Burlington, *Mrs. Flynn.*
SILENE DICHOTOMA, Ehrh. Townshend, *W. H. Blanchard.*
DIANTHUS PLUMARIUS, L. Manchester, *Mrs. L. Munson.*
DELPHINIUM AJACIS, L. Middlebury, *Brainerd*; Westminster, *Eggleson.*
PAPAVER INTERMEDIUM, Becker. Townshend, *W. H. Blanchard.*
P. RHOEAS, L. Townshend, *W. H. Blanchard.*
RAPHANUS SATIVUS, L. Hartland, *Eggleson*; Westminster, *W. H. Blanchard.*
CICER ARIETINUM, L. Burlington, *Mrs. Flynn.*
VICIA FABA, L. Peacham, *F. Blanchard.*
ERODIUM BOTRYS, Bertol. Gravelly bed of Winooski River below woolen mills, Burlington, *Brainerd.*
Gilia linearis, Gray. Burlington, *Mrs. Flynn.*
AJUGA REPTANS, L. Lawn, Burlington, *Jones.*
SHERARDIA ARVENSIS, L. Introduced with seed of Eschscholtzia, Middlebury, *Brainerd.*
AMBROSIA TRIFIDA, L. var. integrifolia, T. & G. Burlington, *Mrs. Flynn.*
ANTHEMIS TINCTORIA, L. Townshend, *W. H. Blanchard.*
ARTEMISIA GNAPHALODES, Muhl. Along railroad, Bellows Falls, *W. H. Blanchard.*
CREPIS SETOSA, Haller. Townshend, *W. H. Blanchard.*
CREPIS VIRENS, var. AGRESTIS, Koch. Townshend, *W. H. Blanchard.*
CREPIS VIRENS, var. RUNCINATA, Bischoff. Townshend, *W. H. Blanchard*; Stowe, *Mrs. H. E. Straw.*
DYSODIA CHRYSANTHEMOIDES, Lag. (Boebera papposa, Rydb.) In river gravel below Winooski Falls, Burlington, *Brainerd.*
HIERACIUM PRAEALTUM, Vill. Burlington, *Mrs. Flynn.*
IVA XANTHIIFOLIA, Nutt. Manchester, *Mrs. L. Munson.*
-

PLANTS NEW TO THE FLORA OF NEW BEDFORD.—The following plants additional to those mentioned in RHODORA, vi. 22, 23, were collected by the writer in the summer of 1903:—*Heracleum Spondylium*, L. was found on the side of a wooded lot and near the highway in the vicinity of New Bedford. If I am not mistaken, this is the first report of this European species from New England. I am indebted to Mr. M. L. Fernald for its identification. *Foeniculum vulgare*, Mill., fennel, was found near the city in waste ground, and afterwards was discovered to be cultivated quite extensively by a Portuguese farmer in the adjoining town of Dartmouth. A considerable patch of *Rubus argutus*, Link, var. *Randii*, Bailey, was also found in the latter town.—E. WILLIAMS HERVEY, New Bedford, Massachusetts.